arXiv:1608.04898v2 [math.RA] 20 Sep 2016

VON NEUMANN FINITE ALTERNATIVE ALGEBRAS

PATRIK NYSTEDT

University West, Department of Engineering Science, SE-46186 Trollhättan, Sweden

Abstract. We show that alternative involutive algebras are von Neumann finite. We use this to show that finite-dimensional alternative algebras are von Neumann Finite. We also show that all the algebras in the infinite chain of classical Cayley-Dickson doublings, starting with the real numbers, are von Neumann finite

1. Introduction

A unital ring A is called *von Neumann finite* (or Dedekind finite, or weakly 1-finite, or affine finite, or directly finite, or inverse symmetric) if every one-sided inverse in A also is two-sided, in other words, if for all $a, b \in A$ satisfying $ab = 1$, the relation $ba = 1$ also holds. Many different classes of *associative* rings have been shown to be von Neumann finite, for instance noetherian, self-injective, PI-rings or reversible rings [5]. Also group rings over fields of zero characteristic [3, 6, 7], or, more generally, endomorphism rings induced by permutation modules [8, 9], have been shown to be von Neumann finite. Von Neumann finiteness for group rings of positive characteristic has, however, remained an open problem (for partial results, see [1] and [2]). Of course, not all unital rings are von Neumann finite, for instance if V is a vector space over a field F, then it is easy to see that the endomorphism ring $A = \mathrm{End}_F(V)$ is von Neumann finite if and only if V is finite dimensional.

In this article, we show three results concerning von Neumann finiteness for *non-associative* rings (see Theorem 1, Theorem 2 and Theorem 3). It seems to the author of the present article that this line of investigation has not previously been considered. Before we state our results, we need some notation. For the rest of the article, let A be an algebra over a field F. By this we mean that A is simultaneously a vector space over F and a, not necessarily associative, ring such that the ring multiplication is F-bilinear.

E-mail address: patrik.nystedt@hv.se.

2010 *Mathematics Subject Classification.* 17D05, 16K20, 17C55, 16W10, 16U99.

Key words and phrases. von Neumann finite, Dedekind finite, directly finite, alternative algebra, involutive algebra.

We always assume that A is unital and that F is contained in the center $Z(A) = \{a \in A \mid a$ commutes and associates with all elements of $A\}$ of A. We let the common identity element of F and A be denoted by 1. Recall that A is called *alternative* if for all $a, b \in A$, the relations $a(ab) = a^2b$ and $(ab)b = ab^2$ hold. We say that A is *involutive* if there is an F-linear operator $A \ni a \mapsto \overline{a} \in A$ satisfying $\overline{ab} = \overline{b}\overline{a}$, $\overline{\overline{a}} = a$, $t(a) := a + \overline{a} \in F$ and $n(a) := a\overline{a} \in F$ for all $a, b \in A$.

In Section 2, we show the following result.

Theorem 1. *If A is alternative and involutive, then A is von Neumann finite.*

In Section 3, we use Theorem 1 to show that Cayley algebras (see Example 8) are von Neumann finite (see Proposition 9). This is in turn used to show the following result.

Theorem 2. *If A is finite dimensional and alternative, then A is von Neumann finite.*

Starting with the real numbers $A_1 = \mathbb{R}$, equipped with the trivial involution, the Cayley-Dickson double process (see Example 8), with $\mu = -1$ at each step, successively yields the complex numbers $A_2 = \mathbb{C}$, Hamilton's quaternions $A_3 = \mathbb{H}$, Graves' octonions $A_4 = \mathbb{O}$, the sedenions $A_5 = \mathbb{S}$, the trigintaduonions $A_6 = \mathbb{T}$, the sexagintaquatronions A_7, the centumduodetrigintanions A_8, the ducentiquinquagintasexions A_9 etc. Their properties keep degrading. Namely, $\mathbb{C}$ is not ordered, $\mathbb{H}$ is not commutative, $\mathbb{O}$ is not associative and neither $\mathbb{S}$, nor any of it's doublings, is alternative. However, all algebras A_i, for $i \geq 1$, keep the property of being *flexible* i.e. for all $a, b \in A_i$ the relation $a(ba) = (ab)a$ holds (see e.g. [10]). In Section 4, we show that these algebras also retain the property of being von Neumann finite.

Theorem 3. *All the algebras A_i, for $i \geq 1$, are von Neumann finite.*

Acknowledgement. The author is indebted to J. Bales for a useful discussion concerning some of the results in Section 4.

2. Involutive Alternative Algebras

In this section, we prove Theorem 1. To this end, we recall some fairly well known facts (see Propositions 4-7). For the convenience of the reader, we include proofs of all of these results (except for Proposition 6(a)). At the end of this section, we use Theorem 1 to show that Cayley algebras (see Example 8) are von Neumann finite (see Proposition 9).

Proposition 4. *If A is involutive, then $\overline{1} = 1$ and $n(1) = 1$.*

Proof. $\overline{1} = 1\overline{1} = \overline{\overline{1}}\,\overline{1} = \overline{1\overline{1}} = \overline{\overline{1}} = 1$ and $n(1) = 1\overline{1} = \overline{1} = 1$. □

Proposition 5. *If A is involutive, then $n(a) = n(\overline{a})$, for $a \in A$.*

Proof. Take $a \in A$. Then $n(\overline{a}) = \overline{a}\,\overline{\overline{a}} = \overline{a}a = \overline{a}a + \overline{a}^2 - \overline{a}^2 = \overline{a}(a + \overline{a}) - \overline{a}^2 = \overline{a}t(a) - \overline{a}^2 = \overline{at(a) - a^2} = \overline{a(a + \overline{a}) - a^2} = \overline{a^2 + a\overline{a} - a^2} = \overline{a\overline{a}} = \overline{\overline{a}}\,\overline{a} = a\overline{a} = n(a)$. □

In the next proposition, we will use the following notation. If $a, b, c \ldots \in A$, then $\langle a, b, c \ldots \rangle$ denotes the unital subalgebra of A generated by $a, b, c \ldots$

Proposition 6. *Suppose that $a, b \in A$.*

(a) *If A is alternative, then $\langle a, b \rangle$ is associative.*
(b) *If A is alternative and involutive, then $\langle a, \overline{a}, b, \overline{b} \rangle$ is associative.*

Proof. (a) See [11, Theorem 3.1].

(b) This follows from (a) and the fact that $F \subseteq Z(A)$ since, for every $a, b \in A$, $\overline{a} = t(a) - a \in F + Fa$ and $\overline{b} = t(b) - b \in F + Fb$, and thus $\langle a, \overline{a}, b, \overline{b} \rangle = \langle a, b \rangle$ is associative. □

Proposition 7. *If A is alternative and involutive, then $n(ab) = n(a)n(b)$, for $a, b \in A$.*

Proof. Take $a, b \in A$. By Proposition 6(b) and the fact that $n(a), n(b) \in F \subseteq Z(A)$, we get that $n(ab) = ab\overline{ab} = ab\overline{b}\overline{a} = an(b)\overline{a} = a\overline{a}n(b) = n(a)n(b)$. □

Proof of Theorem 1. Suppose that $a, b \in A$ satisfy $ab = 1$. We wish to show that $ba = 1$. From Propositions 4-7, it follows that $ba = b1a = b\overline{1}a = b\overline{ab}a = b\overline{b}\,\overline{a}a = n(b)n(\overline{a}) = n(b)n(a) = n(a)n(b) = n(ab) = n(1) = 1$. □

Example 8. Suppose that B is an involutive F-algebra and that μ is a non-zero element of F. Then there is an involution and a multiplication on the F-vector space $A = B \times B$ defined by the relations $\overline{(a, b)} = (\overline{a}, -b)$ and $(a, b)(c, d) = (ac + \mu d\overline{b}, \overline{a}d + cb)$, respectively, for $a, b, c, d \in B$. The resulting structure, called the *Cayley-Dickson double* of B, is an involutive F-algebra with multiplicative identity $(1, 0)$. One can show that A is alternative if and only if B is associative (see [11, p. 46]). If B is a *quaternion algebra*, i.e. a 4-dimensional associative central simple F-algebra, then A is called a *Cayley algebra*. Therefore, Cayley algebras are involutive and alternative. Thus, from Theorem 1, the next result follows immediately.

Proposition 9. *Cayley algebras are von Neumann finite.*

3. Finite Dimensional Alternative Algebras

In this section, we prove Theorem 2. To this end, we first prove two results concerning quotients of algebras and associative algebras (see Proposition 10 and Proposition 11 respectively). Recall that an ideal I of an alternative algebra A is called a *nilideal* if each of its elements is nilpotent. Note that from Proposition 6(a) it follows that powers of elements (and hence also nilpotent elements) can be unambiguously defined for such algebras.

Proposition 10. *If A is alternative and I is a nilideal of A, then A is von Neumann finite if and only if A/I is von Neumann finite.*

Proof. The "only if" statement is immediate. Now we show the "if" statement. Suppose that A/I is von Neumann finite. Take $a, b \in A$ such that $ab = 1$. We wish to show that $ba = 1$. Since A/I is von Neumann finite, there is $c \in I$ such that $ba = 1 + c$. Since I is nilpotent, there is a positive integer n such that $c^n = 0$. We claim that for every positive integer m, the relation

$$(ba - 1)^m = (-1)^m(1 - ba) \tag{1}$$

holds. If we assume that this claim holds, then using (1), with $m = 2n + 1$, we get $ba = ba+0 = ba+c^{2n+1} = ba+(ba-1)^{2n+1} = ba+(-1)^{2n+1}(ba-1) = ba - (ba - 1) = ba - ba + 1 = 1$. Now we show the claim using induction over m.

Base case: $m = 1$. This is clear since $ba - 1 = (-1)(1 - ba)$.

Induction step: suppose that $(ba - 1)^m = (-1)^m(1 - ba)$ holds for some positive integer m. Then, from the induction hypothesis, it follows that $(ba - 1)^{m+1} = (ba - 1)^m(ba - 1) = (-1)^m(1 - ba)(ba - 1) = (-1)^m(ba - 1 - baba + ba) = (-1)^m(ba - 1 - b1a + ba) = (-1)^m(ba - 1 - ba + ba) = (-1)^m(ba - 1) = (-1)^{m+1}(1 - ba)$. □

Proposition 11. *If A is associative and finite dimensional, then A is von Neumann finite.*

Proof. This follows immediately from the fact that such algebras are noetherian rings. Here we give a direct argument. Supppose that $a, b \in A$ are chosen so that $ab = 1$. We wish to show that $ba = 1$. It is clear that $Aa^m \supseteq Aa^{m+1}$ for all positive integers m. Thus, finite dimensionality of A implies that $Aa^n = Aa^{n+1}$ for some positive integer n. Hence, there is $c \in A$ such that $a^n = ca^{n+1}$. From associativity of A it follows that $1 = 1^n = (ab)^n = a^nb^n = ca^{n+1}b^n = caa^nb^n = ca(ab)^n = ca1^n = ca$. Therefore, $b = 1b = cab = c1 = c$ and hence $ba = ca = 1$. □

Proof of Theorem 2. Suppose that A finite dimensional and alternative. We wish to show that A is von Neumann finite. There is a unique maximal nilideal I of A (see [11, Theorem 3.2] and the discussion thereafter). By a Theorem of Zorn (see [11, Theorem 3.12]) A/I is semisimple, that is A/I is the finite direct sum of simple alternative F-algebras. But from another result by Zorn (see [11, Theorem 3.17]) it follows that each such summand either is a finite dimensional associative algebra or a Cayley algebra. Thus, von Neumann finiteness of A follows from Proposition 9, Proposition 10 and Proposition 11. □

4. A Class of Non-alternative Algebras

In this section, we show Theorem 3. To this end, we show three results concerning reversible involutive rings (see Proposition 12, Proposition 13 and Proposition 14). Recall that A is called *reversible* if for all $a, b \in A$ satisfying $ab = 0$, the relation $ba = 0$ also holds. We say that A is *anisotropic* if for

every $a \in A$, the relation $n(a) = 0$ implies that $a = 0$. We say that $n(A)$ has *square roots* if for every $a \in A$, there is $f \in F$ such that $f^2 = n(a)$. We say that A is *real* if $F = \mathbb{R}$.

Proposition 12. *If A is reversible and involutive, then, for all $a, b \in A$, with $n(a) = 1$ (or $n(b) = 1$) and $ab = 1$, the relation $ba = 1$ also holds.*

Proof. Take $a, b \in A$ with $n(a) = 1$ and $ab = 1$. We wish to show that $ba = 1$. Put $c = b - \overline{a}$. Then $ac = a(b - \overline{a}) = ab - a\overline{a} = 1 - n(a) = 1 - 1 = 0$. Since A is reversible, we get that $ca = 0$. Thus $(b - \overline{a})a = 0$ and hence $ba - \overline{a}a = 0$. From Proposition 5, we now get that $ba = \overline{a}a = \overline{a}\,\overline{\overline{a}} = n(\overline{a}) = n(a) = 1$. The case when $n(b) = 1$ is treated similarly. □

Proposition 13. *If A is reversible, involutive, anisotropic and $n(A)$ has square roots, then A is von Neumann finite.*

Proof. Take $a, b \in A$ with $ab = 1$. We wish to show that $ba = 1$. Since A is anisotropic and $n(A)$ has square roots, there is a non-zero $f \in F$ with $f^2 = n(a)$. Put $c = f^{-1}a$ and $d = fb$. Then $n(c) = f^{-1}a\overline{f^{-1}a} = f^{-2}a\overline{a} = f^{-2}n(a) = 1$ and $cd = f^{-1}afb = ff^{-1}ab = 1$. From Proposition 12, we thus get that $dc = 1$. Hence $1 = fbf^{-1}a = ff^{-1}ba = ba$. □

Proposition 14. *Suppose that A is involutive and real. If for all non-zero $a, b \in A$, the relations $n(a) > 0$ and $n(ab) = n(ba)$ hold, then A is von Neumann finite.*

Proof. From Proposition 13, it follows that it is enough to show that A is reversible, anisotropic and $n(A)$ has square roots. Since $n(a) > 0$, for non-zero $a \in A$, it is clear that A is anisotropic and $n(A)$ has square roots. Now we show that A is reversible. Take $a, b \in A$ with $ab = 0$. Then $n(ba) = n(ab) = n(0) = 0$. Since A is anisotropic, we get that $ba = 0$. □

Proof of Theorem 3. We know from Example 8 that each A_i is involutive. Also, from [4, Lemma 1.4.2], we know that each A_i has the property that $n(ab) = n(ba)$, for $a, b \in A_i$. Thus, from Proposition 13, it follows that we need to prove the following statement for all $i \geq 1$: for every non-zero $a \in A_i$, $n(a) > 0$. We will prove this by induction over i. Base case: $i = 1$. This is clear. Induction step: suppose that the statement holds for some $i \geq 1$. Take $a, b \in A_i$ with a or b non-zero. Then, from the induction hypothesis, it follows that

$$n((a,b)) = (a,b)\overline{(a,b)} = (a,b)(\overline{a}, -b) = (a\overline{a} - (-b)\overline{b}, \overline{a}(-b) + \overline{a}b) =$$

$$= (a\overline{a} + b\overline{b}, 0) = (n(a) + n(b), 0) = n(a) + n(b) > 0.$$

Thus, the statement also holds for A_{i+1}. □

References

[1] P. Ara, K. OMeara, and F. Perera. Stable finiteness of group rings in arbitrary characteristic. *Adv. Math.* **170**, 224-238 (2002).

[2] G. Elek and E. Szab. Sofic groups and direct finiteness. *J. Algebra* **280**, 426-434 (2004).
[3] I. Kaplansky. *Fields and Rings*. Univ. of Chicago Press, Chicago, Ill. (1969).
[4] S. H. Khalil and P. Yiu. The Cayley-Dickson algebras, a theorem of Hurwitz, and quaternions, *Bulletin de la societe des sciences et des lettres de Lod z*, vol. XLVII. 117–169 (1997)
[5] T. Y. Lam. *Exercises in Modules and Rings*. Springer-Verlag (2007).
[6] S. Montgomery. Left and Right Inverses in Group Algebras. *Bull. AMS* **75** ,539–540 (1969).
[7] D. S. Passman, Idempotents in group rings. *Proc. Amer. Math. Soc.* **28**, 371–374 (1971).
[8] P. Lundström. Von Neumann Finite Endomorphism Rings. *Indag. Mathem.*, N.S., **14** (2), 223–232 (2003).
[9] P. Lundström. Brauer Group Rings. *JP Journal of Algebra, Number Theory & Applications*, **4**, 465–494 (2004).
[10] R.D. Schafer. On the algebras formed by the Cayley-Dickson process, *Amer. J. Math.* **76** 435–446 (1954).
[11] R. D. Schafer. *An Introduction to Nonassociative Algebras*. Dover (1995).